



LipoReM

Premium Lipolysis Solution

Serum for the Slim

This is a liquid solution which is used in renewal of your local fatty volume and for people which have excessive fat due to aging and high fat diet.

Slim line is formed as making your skin look firm and smooth.

TYPE

Liquid

VOLUME

10ml x 5 vials

EXPIRY DATE

2 years from date of manufacture

(This is single use. Use all after opening)

STORAGE

Store at room temperature

APPLICATION PART

Chin, Abdomen, Love handles, Upper arms, Thigh

LipoReM Information

Ingredients	Tissue reliever, saline solution
Usage	LipoReM is used after absorbing the solution in the container using a syringe of 10~30ml.
Storage	Storage at room temperature within the expiration date (Do not store remaining solution after use)

Injection site	All areas with fat
Injection point	1~1.5cm interval
Injection volume per point	Face : 0.05~0.1ml, Body : 0.1~0.3ml

Application area		Usage*	Notes
Face	V-line, Double chin	1.0-1.5ml	Injection Dosage: - Do not exceed total 4 ml for face for 30 days per person. - Maximum usage (dosage) of each part of the face for single treatment should not exceed the noted dosage stated in the Table. - The same site can be treated every 30days. Injection Technique: - Inject depth 5mm, injection volume per point is 0.05ml ~ 0.1ml, injection interval is 1~1.5cm.
	Cheek, Bucal fat	0.5-1.0ml	
	Below ear	0.5-1.0ml	
	Malar	0.5ml	
Body	Flank	15-20ml	Injection Dosage: - Do not exceed total 30ml for body for 30 days per person. - Maximum usage (dosage) of each part of the body for single treatment should not exceed the noted Dosage stated in the Table. - Consume a total 1-3 vials (each vial 10ml) of LipoReM per treatment for body depending on the treatment area. It is recommended to repeat this treatment 2-3 times at 15 days interval which will lead a patient to consume 1-2 boxes to complete a whole process of treatment. - The same site can be treated every 30days. Injection Technique: - Injection volume per point is 0.1ml~0.3ml, injection interval is 1~1.5cm.
	Belly	10ml	
	Bra-line	20ml	
	Thigh	20ml	
	Calf	10-15ml	
	Arm	8-10ml	
	Ankle	6ml	
Trapezius line	15-20ml		

*Usage - Maximum dosage for one area of the body for per treatment.
While the maximum Injection Volume per point should be considered.

Efficacy	7-30 days
Precaution	Facial wash, shower is recommended after 6 hours
	Individually swelling and bruising may occur. Massage after the application is recommended
	When using more than 20ml per day, take antibiotics for 4 days (3 times a day)
Advantage	Volume reduction is fast
	No side effects
	Immediately life is available
	Simple therapy
Disadvantage	The large area should be divided at interval (2-3 times)

Treatment Area



✓ Forearm



✓ Double chin



✓ Side belly




✓ Belly fat


Treatment Method



1
Shake LipoReM vials well to mix the deposited particles on the bottom and open the plastic cap.



2
Insert the needle into the rubber top of the vial, pull back the plunger to the line on your syringe for your dose.



3
Remove the needle from the vial. The medicine is ready to be injected into the treatment area.

Before & After

✓ Double Chin Treatment



LipoReM

PREMIUM LIPOLYSIS SOLUTION

Caution

- Avoid stimulation, make-up or cleansing on treatment area for 6 hours after treatment and gently cleanse on the day of treatment.
- Effect of treatment will appear within 1 week after treatment. Occasionally, people with large or thick layer of fat may experience slower effect within about 1 or 2 weeks.
- It is recommended to avoid intense sports, sauna, drinking and smoking for about 1 week after treatment (Alcohol may interrupt breakdown of fat that the effect of treatment may be insignificant or none).
- Normally, there are no side-effects after the treatment but occasionally, treatment area may be slightly swollen or bruised.
- Depending on physical constitution, allergic reactions may occur.
- Product includes small amount of steroids that irregular menstruation may occur for women.
- As the results of treatment may vary depending on individuals, additional treatments of 1 to 3 times may occur for face and 3 to 5 times for body.
- It is recommended to have additional treatments with 4 weeks interval.
- Prescribed antibiotic medicines help reducing swelling and allergic reactions and are recommended to take.
- After body treatment, It is advised either to massage with oil or frequently apply body lotion to the treatment area, as skin may be dry during the hours of that breakdown